

Organic Acid Tests (OATs)

TEST	RESULT	UNITS	NORMAL RANGE	
Nutritional Organic Acids (Urine)				
Vitamin B12 Marker - May be deficient if high				
Methylmalonate (MMA)	Above range	4.9	ug/mg	0 - 2.5
Vitamin B6 Markers - May be deficient if high				
Xanthurenate	Above range	1.23	ug/mg	0.12 - 1.2
Kynurenate	Above range	5.4	ug/mg	0.8 - 4.5
Biotin Marker - May be deficient if high				
b-Hydroxyisovalerate	Within range	7.9	ug/mg	0 - 12.5
Glutathione Marker - May be deficient if high				
Pyroglutamate	Within range	42.0	ug/mg	28 - 58
Gut Marker - Potential gut putrefaction or dysbiosis if high				
Indican	Above range	114.0	ug/mg	0 - 100
Neuro-Related Markers (Urine)				
Dopamine Metabolite				
Homovanillate (HVA)	Within range	4.4	ug/mg	3 - 11
Norepinephrine/Epinephrine Metabolite				
Vanilmandelate (VMA)	Within range	4.3	ug/mg	2.2 - 5.5
Neuroinflammation Marker				
Quinolinate	Within range	8.0	ug/mg	0 - 9.6
Additional Markers (Urine)				
Melatonin - Waking				
6-OH-Melatonin-Sulfate	Below range	5.3	ng/mg	10 - 85
Oxidative Stress / DNA Damage				
8-Hydroxy-2-deoxyguanosine (8-OHdG)	Within range	2.6	ng/mg	0 - 5.2

- The MMA is above the range. This may indicate vitamin B12 or adenosylcobalamin deficiency. B12 is important for phase 2 methylation (estrogen detox), neurotransmitter synthesis, and other key processes.
- Both the xanthurenate and kynurenate are above the range. This may indicate vitamin B6 deficiency. B6 is important for phase 2 methylation (estrogen detox), neurotransmitter synthesis, and other key processes. Tryptophan taken within 72 hours before testing can also raise these markers without indicating a true B6 deficiency.
- The indican is above the range. This can indicate gut dysbiosis. Gut dysbiosis can affect estrogen metabolism, inflammation, and malabsorption of nutrients. Further GI testing may be indicated.
- The waking urinary 6-OH-Melatonin-Sulfate is low. This reflects low overnight production of melatonin. This may be implicated in poor sleep and insomnia.

Thank you for choosing DUTCH for your functional endocrinology testing needs!

Please review our DUTCH resources for information on reading the DUTCH test:

For DUTCH Overviews and Tutorials, click here: <https://dutchtest.com/tutorials>

To view the steroid pathway chart, click here: <https://dutchtest.com/steroid-pathway>

Finally, please review the patient's results along with their requisition form. It is designed to capture relevant medications, symptoms, diagnoses, sample collection, and notes that may be helpful in interpreting the results.

Additional Comments

Ordering Provider:

Test Provider MD

Male Sample Report

123 A Street

Sometown, CA 90266

DOB: 1976-01-01

Age: 49

Sex: Male

Collection Dates:

2025-06-13 (U1)

Organic Acid Tests (OATs)

TEST		RESULT	UNITS	NORMAL RANGE
Nutritional Organic Acids (Urine)				
Vitamin B12 Marker - May be deficient if high				
Methylmalonate (MMA)	Within range	2.9	ug/mg	0 - 3.5
Vitamin B6 Markers - May be deficient if high				
Xanthurenate	Above range	2.20	ug/mg	0.2 - 1.9
Kynurenate	High end of range	6.4	ug/mg	1 - 6.6
Biotin Marker - May be deficient if high				
b-Hydroxyisovalerate	Above range	23.0	ug/mg	0 - 18
Glutathione Marker - May be deficient if high				
Pyroglutamate	Within range	62.0	ug/mg	38 - 83
Gut Marker - Potential gut putrefaction or dysbiosis if high				
Indican	Within range	90.4	ug/mg	0 - 131
Neuro-Related Markers (Urine)				
Dopamine Metabolite				
Homovanillate (HVA)	Low end of range	4.4	ug/mg	4 - 16
Norepinephrine/Epinephrine Metabolite				
Vanilmandelate (VMA)	High end of range	7.3	ug/mg	2.5 - 7.5
Neuroinflammation Marker				
Quinolate	Within range	9.1	ug/mg	0 - 12.5
Additional Markers (Urine)				
Melatonin - Waking				
6-OH-Melatonin-Sulfate	Below range	7.8	ng/mg	10 - 85
Oxidative Stress / DNA Damage				
8-Hydroxy-2-deoxyguanosine (8-OHdG)	Within range	3.7	ng/mg	0 - 8.8

- The xanthurenate is above the range. This may indicate a vitamin B6 deficiency. B6 is essential for phase 2 methylation (estrogen detoxification), neurotransmitter synthesis, and other key metabolic processes. Tryptophan taken within 72 hours before testing can also raise kynurenate without indicating a true B6 deficiency.
- The b-hydroxyisovalerate is above the range. This may indicate biotin (vitamin B7) deficiency.
- The waking urinary 6-OH-Melatonin-Sulfate is low. This reflects low overnight production of melatonin. This may be implicated in poor sleep and insomnia.

Thank you for choosing DUTCH for your functional endocrinology testing needs!

Please review our DUTCH resources for information on reading the DUTCH test:

For DUTCH Overviews and Tutorials, click here: <https://dutchtest.com/tutorials>

To view the steroid pathway chart, click here: <https://dutchtest.com/steroid-pathway>

Finally, please review the patient's results along with their requisition form. It is designed to capture relevant medications, symptoms, diagnoses, sample collection, and notes that may be helpful in interpreting the results.

Additional Comments